

Before manure is applied, the nutrient requirement of the crop should be determined. Soil testing is a reliable method to determine nutrient levels in the soil. With this information, and knowledge of the nutrient levels of the manure, economically and environmentally sound application rates for both manure and fertilizer can be determined.

Manure application rates are usually based on crop nitrogen requirements. Manure application rates should never exceed the crop nitrogen requirement. However, after long term application, the phosphorus level in the soil will probably build up to a level that will prohibit continued manure applications. All manure application sites should be monitored with a soil testing program. To prevent the build up of soil phosphorus and potassium levels to a very high range, it would be best to calculate a manure application rate on basis of crop phosphorus or potassium removal (do not exceed crop N requirements). [Table 1](#) provides a list of estimated nutrient removal per unit of yield for many crops. Actual removal will be site specific and may be determined by a laboratory analysis of a representative sample of the plant material removed from the field.

NUTRIENT AVAILABILITY

The nutrients contained in animal manures, composts, or other organic materials are less readily available to plants than the nutrients of most inorganic fertilizers. The nutrient content of these materials is highly variable ([Table 2](#)) and a laboratory analysis should be used to determine the actual nutrient content of manure materials.

NITROGEN:

Manure would be classified as a slow release nitrogen fertilizer. Approximate manure nitrogen availability percentages are listed in [Table 3](#). They vary according to the type of material, storage, and application method. When incorporated, most of the ammonium nitrogen is available during the first year. Nitrogen availability percentages listed in [Table 3](#) are total N and account for ammonium nitrogen content. The figures in [Table 3](#) are based on availability of total N during the first year following a spring application.

ORGANIC N:

(Total N minus ammonium N, or assume 60 to 75 percent of the Total N as organic) will supply nitrogen to the cropping system for several years. To estimate residual N use the availability factors listed in [Table 4](#).

PHOSPHORUS AND POTASSIUM:

About 70 percent of the total P applied in manure will become available in the year of application. Of the total potassium, 100 percent will be available the year of application.

DETERMINING NUTRIENT AND FERTILIZER NEEDS

Nitrogen is usually selected as the priority nutrient and total crop nutrients are determined by soil test and crop removal. When soil test results are high or very high in phosphorus or potassium it may be best to use these nutrients to determine the manure application rate. Also, if the manure tests low in nitrogen, basing the application rate on phosphorus or potassium may be best. In any situation, manure application rate should never exceed the crop's nitrogen requirement.

Amounts of nutrients that can be added from manure without greatly increasing soil test levels are estimated on the basis of expected crop removal. Values in [Table 1](#) may be used to estimate crop removal expected per unit of yield for various crops.

Manure sampling, manure analysis, and spreader calibration are part of a comprehensive nutrient management plan. Manure with greater than 20 percent solids is classified as dry manure and is handled as a solid. Manure with 4 to 20 percent solids is classified as semi-solid and can be handled as a liquid. Semi-solid manure usually requires thorough agitation before pumping and sampling. Manure with less than 4 percent solids is classified a liquid manure and is handled with pumps, tank wagons, and irrigation equipment.

A representative sample is needed to provide an accurate manure analysis. One of the many factors affecting the nutrient content of manure is how the manure is handled and stored. Each handling system results in different types of nutrient losses. The most important thing in collecting a manure sample is to obtain it in a similar way to the methods used in developing the standard nutrient values.

WHEN TO SAMPLE MANURE

Sample manure at the time of land application or as close to the time of application as possible. Sampling at the time of application will not provide manure recommendations that can be used to adjust the amount of manure applied. However, the results can be used to adjust future manure applications and to adjust the amount of inorganic fertilizer applied. If you apply manure several times a year, sample when you apply the bulk of the manure.

Ideally, manure sampling should be done in the field as manure is applied. This ensures that losses that occur during handling, storage, and application are taken into account.

MANURE SAMPLING IN THE FIELD

Dry or Solid Field Sampling

To sample manure from barns, holding areas, dry stacks, or feed lots, collect a sample as follows:

Use the "hand and bag" method to collect all solid manure samples. Place a one-gallon resealable freezer bag turned inside out over one hand. Grab a handful of manure with the covered hand and turn the freezer bag right side out over the sample with the free hand. Seal the bag and place it in another freezer bag to prevent leaks. Label the bag and send to the lab or freeze it immediately to prevent nutrient losses. Take three samples for dry or solid manure. Combine the samples and mix. Place in a zip-lock bag.

Liquid Manure Sampling

When sampling liquid manure agitate the manure in the storage facility to obtain a representative sample for laboratory analysis.

Liquid Manure Applied with Spreaders

1. Immediately after filling the tank spreader, use a clean plastic bucket to collect manure from the unloading port or the opening near the bottom of the tank. Be sure the opening does not have solids accumulated that can contaminate the samples.
2. Stir the manure in the pail and immediately fill a one-quart flexible plastic bottle about 25 per-cent full. **Do not use a glass bottle as it might explode from pressure build up.** Squeeze as much air out of the bottle as possible before capping.
3. Put your name, date and sample number on the bottle and the information sheet.
4. If the sample cannot be sent to the laboratory within a few hours, it should be refrigerated. Place the sample in a plastic bag, seal the bag, and keep cool until it is sent to the laboratory. Ship so that the sample arrives promptly at the laboratory.

Liquid Manure Applied by Irrigation Systems

1. Place catch pans or buckets randomly in the field to collect the liquid manure that is applied by an irrigation system.

2. Immediately after the manure has been applied, collect the manure from each pan or bucket and combine in one bucket to make a composite sample.

Mix the manure and fill a one-quart flexible plastic bottle about 25 percent full. Seal and label the bottle and seal in a plastic bag. If the sample cannot be shipped to the laboratory right away, keep refrigerated. Ship to arrive promptly in the laboratory.

Dry or Solid Manure

Paved Lots

1. Collect manure by scraping a shovel across 25 feet of the paved feedlot. Repeat this process six to eight times. Avoid samples from areas that are very wet or contain large amounts of feed and hay.
2. Use the shovel to thoroughly mix manure by scooping the outside of the pile to the center of the pile.
3. Collect a sample using the "hand and bag" method that was de-scribed in the section on dry or solid field sampling.

Barn Gutter

1. Shovel a manure sample to the depth of the gutter from the gutter.
2. Remove the manure from the gutter and place it on the barn floor. Mix the sample by hand (wearing freezer bags) with a kneading motion. When collecting samples from a gutter, be sure to include the liquid that is in the bottom of the gutter.
3. Collect a sample using the "hand and bag" method.
4. Repeat steps one through three from other locations in the gutter to collect three subsamples. Combine the subsamples and mix. Place in zip-lock bag and squeeze out all of the air before closing.

Dry Stack

This is manure stored outside in a stacking shed or above ground solid waste storage facility.

1. Using a pitchfork or shovel, take manure from several locations throughout the dry stack and place it in a pile. Collect samples from the outside and center of the stack.
2. Mix the manure with a shovel by scooping the outside of the pile to the center of the pile.
3. Collect a manure sample by the "hand and bag" method.
4. Repeat steps one through three to collect the three subsamples. Combine the subsamples and mix. Place in a zip-lock bag and squeeze out all of the air before closing.

Table 1. Nutrient Removal by Crops				
		Approximate Pounds per Acre of Nutrients		
		Removed by the Portion of the Crop Shown		
Crop	Unit	N	P₂O₅	K₂O
Corn		1.0	0.37	0.26
Grain	bu.	0.75	0.15	1.06
Stover		1.75	0.52	1.32
Total Removed				
Cotton				
Lint and seeds	lb.	0.08	0.04	0.03
Stalk, etc.		0.06	0.02	0.05
Total Removed		0.14	0.06	0.08
Barley				
Grain	bu.	1.0	0.4	0.3
Straw		0.4	0.1	1.1
Total Removed		1.4	0.5	1.4
Oats				
Grain	bu.	0.7	0.25	0.2

Straw		0.3	0.15	1.25
Total Removed		1.0	0.40	1.45
Peanuts				
Nuts	lb.	0.03	0.01	0.01
Vines		0.02	0.01	0.03
Total Removed		0.05	0.02	0.04
Rice				
Grain	bu.	0.5	0.24	0.14
Straw		0.3	0.11	0.85
Total Removed		0.8	0.35	0.99
Grain Sorghum				
Grain	bu.	0.83	0.41	0.21
Stover		0.94	0.18	1.06
Total Removed		1.77	0.59	1.27
Soybeans				
Beans	bu.	4.00	0.80	1.40
Stover		1.15	0.27	0.96
Total Removed		5.15	1.07	2.36
Tobacco				
Leaf	lb.	0.03	0.01	0.05
Stalk		0.01	0.01	0.03
Total Removed		0.04	0.02	0.08
Wheat				
Grain	bu.	1.25	0.62	0.37
Straw		0.57	0.15	0.90
Total Removed		1.82	0.77	1.27

Table 1. Continued - Nutrient Removal by Crops

			Approximate Pounds per Acre of Nutrients		
			Removed by the Portion of the Crop Shown		
Crop	Unit	Dry Matter %	N	P ₂ O ₅	K ₂ O
Hay					
Alfalfa (early bloom)	Ton	89.2	52	9	43
		89.2	41	9	37
Alfalfa (full bloom)		89.0	40	9	23
Coastal Bermuda		89.0	28	10	30
		91.2	67	16	47
Ryegrass		87.7	42	10	35
Ladino Clover		88.5	28	10	34
Red Clover		93.2	40	10	23
Fescue		88.2	26	8	30
Lespedeza		88.3	31	11	44
Oats		88.9	31	8	32
Orchardgrass		87.6	39	9	20
Sudangrass		88.4	27	7	34
Soybeans		88.2	54	10	44
Timothy		85.9	25	7	31
Vetch		90.6	32	7	30
Wheat					
Peanut Silage					
Alfalfa	Ton	28.0	15	4	17

Barley		35.0	12	4	14
Corn		35.0	8	3	8
Grass-Legume		45.0	16	5	16
Oats		35.0	11	4	15
Grain Sorghum		30.0	8	2	9
Forage Sorghum		30.0	7	2	7
Sudangrass		35.0	10	3	14
Soybeans		30.0	17	7	7
Timothy		37.5	11	3	14
Wheat		35.0	10	3	10
Rye		33.0	12	3	14

**Table 2. Approximate Manure Nutrients
Remaining at Time of Application**

			Lb/Ton			Lb/1000 gal		
Species	System	Solids %	N	P ₂ O ₅	K ₂ O	N	P205	K20
Cattle	Daily Spread	15	8	5	10	---	---	---
	Anaerobic Pit	8	---	---	---	24	18	29
	Earthen Storage	10	---	---	---	32	14	28
	Anaerobic Lagoon	1	---	---	---	4	5	5
	Above Ground Storage	12	---	---	---	46	18	40
	Covered Stack	18	10	9	12	---	---	---

	Anaerobic Pit	4	---	---	---	36	27	19
	Anaerobic Lagoon	1	---	---	---	4	2	4
	Open Feedlot	15	10	7	10	---	---	---
	Liquid Pit	13	---	---	---	80	37	96
	Dry Pit (dry)	85	100	70	40	---	---	---
	Dry Pit (crumbly)	70	60	55	30	---	---	---
	Dry Pit (moist)	50	40	40	20	---	---	---
	Dry Pit (fresh)	25	30	20	10	---	---	---
	Compost	54	44	66	48	---	---	---
	Dry Pit	25	23	8	20	---	---	---
	Dry Pit	20	12	5	9	---	---	---
	Dry Pit	54	44	66	48	---	---	---

Table 3. Nitrogen Availability Coefficients for Manure Total Nitrogen Content

Type of Manure	Method of Incorporation			
	Broadcast, No Incorporation	Broadcast, Incorporation within 12 hours	Injected	Irrigated, No Incorporation
Dairy (semi-solid)	0.40	0.60	---	---
Dairy (slurry)	0.45	0.60	0.70	0.45
Beef	0.40	0.60	---	---
Swine (slurry)	0.40	0.60	0.70	0.40
Swine Lagoon	0.50	0.80	0.85	0.50
Poultry	0.50	0.50	0.50	---
Municipal sludge	0.50	0.50	0.50	---
Composted Materials	0.20	0.20	---	---

**Table 4. Residual Availability of Organic N
in Manure and Compost**

Material Applied	Organic N Availability Coefficients
Last Year	0.13
2 Years Ago	0.05
3 Years Ago	0.02